

## RESEARCH ARTICLE

# Molecular characterization and biofilm-formation analysis of *Listeria monocytogenes*, *Salmonella* spp., and *Escherichia coli* isolated from Brazilian swine slaughterhouses

Rebecca Lavarini dos Santos<sup>1</sup>\*, Emilia Fernanda Agostinho Davanzo<sup>1</sup>, Joana Marchesini Palma<sup>1</sup>, Virgílio Hipólito de Lemos Castro<sup>1</sup>, Hayanna Maria Boaventura da Costa<sup>1</sup>, Bruno Stéfano Lima Dallago<sup>1</sup>, Simone Perecmanis<sup>1</sup>, Ângela Patrícia Santana<sup>1</sup>

Faculty of Agronomy and Veterinary Medicine, University of Brasília (UnB), Brasília, Federal District, Brazil

\* These authors contributed equally to this work.

\* [rebecca.lavarini@gmail.com](mailto:rebecca.lavarini@gmail.com)



## OPEN ACCESS

**Citation:** dos Santos RL, Davanzo EFA, Palma JM, Castro VHdL, da Costa HMB, Dallago BSL, et al. (2022) Molecular characterization and biofilm-formation analysis of *Listeria monocytogenes*, *Salmonella* spp., and *Escherichia coli* isolated from Brazilian swine slaughterhouses. PLoS ONE 17(9): e0274636. <https://doi.org/10.1371/journal.pone.0274636>

**Editor:** Yung-Fu Chang, Cornell University, UNITED STATES

**Received:** May 3, 2022

**Accepted:** August 31, 2022

**Published:** September 20, 2022

**Copyright:** © 2022 dos Santos et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its [Supporting Information](#) files.

**Funding:** the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (R\$ 36.000) and the DPG of University of Brasília (R\$ 8.000) as funders, both provided financial support for the author Rebecca Lavarini dos Santos to carry out the research. The funders had no role in study

## Abstract

This study aimed to verify the presence of *Listeria monocytogenes*, *Salmonella* spp., and *Escherichia coli* in two Brazilian swine slaughterhouses, as well as to perform antibiograms, detect virulence and antimicrobial resistance genes, and evaluate the *in vitro* biofilm-forming capability of bacterial isolates from these environments. One *Salmonella* Typhi isolate and 21 *E. coli* isolates were detected, while *L. monocytogenes* was not detected. *S. Typhi* was isolated from the carcass cooling chamber's floor, resistant to several antimicrobials, including nalidixic acid, cefazolin, chloramphenicol, doxycycline, streptomycin, gentamicin, tetracycline, and sulfonamide, and contained resistance genes, such as *tet(B)*, *tet(C)*, *tet(M)*, and *ampC*. It also showed moderate biofilm-forming capacity at 37°C after incubating for 72 h. The prevalence of the 21 *E. coli* isolates was also the highest on the carcass cooling chamber floor (three of the four samplings [75%]). The *E. coli* isolates were resistant to 12 of the 13 tested antimicrobials, and none showed sensitivity to chloramphenicol, an antimicrobial prohibited in animal feed since 2003 in Brazil. The resistance genes *MCR-1*, *MCR-3*, *sul1*, *ampC*, *clmA*, *cat1*, *tet(A)*, *tet(B)*, and *blaSHV*, as well as the virulence genes *stx-1*, *hlyA*, *eae*, *tir $\alpha$* , *tir $\beta$* , *tir $\gamma$* , and *saa* were detected in the *E. coli* isolates. Moreover, 5 (23.8%) and 15 (71.4%) *E. coli* isolates presented strong and moderate biofilm-forming capacity, respectively. In general, the biofilm-forming capacity increased after incubating for 72 h at 10°C. The biofilm-forming capacity was the lowest after incubating for 24 h at 37°C. Due to the presence of resistance and virulence genes, multi-antimicrobial resistance, and biofilm-forming capacity, the results of this study suggest a risk to the public health as these pathogens are associated with foodborne diseases, which emphasizes the hazard of resistance gene propagation in the environment.

design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

*Escherichia coli*, *Salmonella* spp., and *Listeria monocytogenes* are among the main bacteria involved in foodborne diseases, and have been evaluated in depth to prevent future outbreaks across the world [1,2].

*Salmonella* spp. is mostly involved in foodborne illnesses worldwide [3]. Approximately 2,500 *Salmonella* serotypes have been identified, the majority of which may adapt to several animal hosts, including humans [4]. According to the Epidemiological Profile of Etiological Agents published by the Brazilian Ministry of Health [5], *E. coli* is the second most common bacterial agent involved in food poisoning outbreaks in Brazil. In addition, this bacterium also causes foodborne outbreaks worldwide and its presence indicates fecal contamination [6,7].

Furthermore, the persistence of foodborne pathogens in biofilms has also been reported, mostly on food contact surfaces, affecting product quality, quantity, and safety [8]. In the meat industry, bacterial biofilms are a major concern due to accumulation in areas difficult to sanitize, leading to cross-contamination and food spoilage [9–11]. In food processing units, *Listeria* spp. has been detected on equipment surfaces, impermeable sealing substances, conveyor belts, and drains, persisting in the industrial environment from months to years [12]. Moreover, *Listeria* spp. can grow at 4–10°C, which is the temperature range commonly used to control food infections, and can become a problem during food handling [13,14].

The presence of these pathogenic microorganisms is a safety hazard to food industries, since they are unlikely to be eliminated from the processing line due to their proliferation and possible biofilm formation [12,15], thus increasing resistance to sanitizers as well as physical and chemical treatments [9,16]. In addition to compromising food hygiene and posing a public health risk, antibiotic resistance and gene transfer among bacteria are associated, potentially increasing the number of circulating virulent strains [17–19].

Since Brazil is the fourth largest pork exporter, and a good performance in this market is due to competitive prices coupled with quality products, it is essential to pay attention to pathogenic microorganisms that can lead to sanitary crises or represent barriers to commercialization [20]. Estimating the number of foodborne outbreaks related to pork meat is difficult due to the lack of reliable data; the contamination rate is under-reported as the majority of cases are not registered [21].

Meanwhile, there have been few reports of biofilms in Brazilian pork industries and there is an absence of data in the Federal District of Brazil and the surrounding region. This study aimed to detect *E. coli*, *Salmonella* spp., and *L. monocytogenes* in the environment and equipment of swine slaughterhouses in the Federal District of Brazil. Molecular characterization and antimicrobial resistance testing of strains isolated from biofilms were also conducted.

## Material and methods

### Origin of the samples

Samples were collected from two swine slaughterhouses (A and B) located in the Federal District of Brazil and two visits were made to each swine slaughterhouse between 2019 and 2021, with a minimum 24 h interval. Swabbing (Absorve®; São Paulo, Brazil) of a delimited area was used for sampling the surfaces, equipment, and utensils [22]. A total of 44 swab samples were collected from 11 points each of two slaughterhouses (A and B) during two visits, using one swab per point per visit. The sample points were defined according to the protocols presented by Cabral *et al.* [23], Nicolau & Bolocan [24], and Barros *et al.* [22] divided into facilities (floors, walls, and drains) and equipment/utensils (tables, bleeding knife, dehairing machine, and carcass splitting saw).

Samples were collected between the last post-slaughter hygiene process at the end of the workday and before starting the daily activities with pre-slaughter hygiene procedures, due to the relation of bacterial permanence on surfaces post hygiene in industries with the presence of bacterial biofilms [25,26].

### ***Salmonella* spp., *L. monocytogenes*, and *E. coli* isolation**

*E. coli* was isolated from the swab samples and identified using a previously described methodology [27]. Briefly, the swabs were transferred from tubes containing 0.1% peptone water (HiMedia®; Mumbai, India) to tubes containing 9 mL 1% buffered peptone water (Acumedia®; Melbourne, Australia) and incubated at 37°C for 24 h. Subsequently, they were streaked onto Eosin Methylene Blue agar plates and incubated at 37°C for 24 h to observe the growth of typical *E. coli* colonies (blue-black colonies with or without metallic green reflex). The *E. coli* colonies were subjected to standard biochemical tests for microbial identification [28,29].

For *L. monocytogenes* isolation, swab samples were analyzed according to the methodology described by the Brazilian Normative Instruction n° 40 [30] for research and microbial *L. monocytogenes* isolation. The surface swabs were transferred from tubes containing 0.1% peptone water to tubes containing 9 mL 1% buffered peptone water and incubated at 37°C for 24 h. After incubation, 1 mL culture was transferred to 9 mL UVM broth (Acumedia®) and incubated at 35°C for 24 h. Then, 0.1 mL culture was transferred to 10 mL Fraser broth (Acumedia®) and incubated at 35°C for 24 h. Fraser broth tubes with observed esculin hydrolysis were plated on MOX agar plates (Difco™; Berkshire, England) and incubated at 35°C for 24 h. Small colonies with a halo of esculin hydrolysis were collected, transferred to Brain Heart Infusion (BHI) broth (Difco™), and incubated at 37°C for 24 h. A turbidity test was performed using droplets obtained from the BHI broth. Gram staining and catalase tests were performed [31].

To identify *Salmonella* spp., swab samples were analyzed according to the protocols described in the Technical Manual for Laboratory Diagnosis of *Salmonella* spp. [32] and ISO 6579/2002 [33]. Briefly, the surface swabs were transferred from tubes containing 0.1% peptone water to tubes containing 9 mL 1% buffered peptone water and incubated at 37°C for 24 h. After incubation, 1 and 0.1 mL cultures were transferred to 10 mL Selenite cystine broth (Merck®; Darmstadt Germany) and Rappaport Vassiliadis broth (Fluka™; Buchs, Germany), respectively, and incubated at 42°C for 24 h. Next, the above-mentioned broths were streaked onto selective modified Brilliant-green Phenol-red Lactose Sucrose agar plates (Acumedia®) and incubated at 37°C for 24 h. Three colonies with morphological characteristics of *Salmonella* spp. were streaked on Triple Sugar Iron (Acumedia®) agar slants and incubated at 37°C for 18–24 h. TSI tubes with potential *Salmonella* growth were biochemically tested as indicated in the Technical Manual for Laboratory Diagnosis of *Salmonella* spp. [32], including urea hydrolysis, phenylalanine deaminase, indole production, Voges–Proskauer test, methyl red test, lysine decarboxylase, and citrate utilization. Positive controls for standardization were provided by the Oswaldo Cruz Foundation, Rio de Janeiro.

Colony PCR analysis [34] was performed to identify and confirm *Salmonella* spp. and amplification reactions were performed in a final volume of 25 µL, containing 2 Units Taq DNA polymerase (Invitrogen®. Waltham, MA, USA), 2 mM phosphate deoxyribonucleotides (Invitrogen®), 1× buffer (200 mM Tris-HCl, pH 8.4; 500 mM KCl; Invitrogen®), 1.5 mM MgCl<sub>2</sub> (Invitrogen®), and 1 µM primers. The reaction was performed in a MyCycler thermal cycler (BioRad®; Hercules, CA, USA) under the following conditions: initial denaturation at 94°C for 3 min, 35 cycles at 94°C for 0.40 min, annealing temperature according to each primer for 1.15 min, and 72°C for 1.15 min; and a final cycle at 72°C for 7 min. The expected

**Table 1. *Salmonella* spp. research detection primers.** Oligonucleotides used for *Salmonella* spp. confirmation and serovar detection of *Salmonella* spp.

Gene	Primer	Oligonucleotide sequence (5'→3')	Size (bp)	Annealing temperature (°C)	Reference
<i>ompC</i>	OMPCF	ATCGCTGACTTATGCAATCG	204	57	[35]
	OMPCR	CGGGTTGCGTTATAGGTCTG			
<i>entF</i>	ENTF	TGTGTTTTATCTGATGCAAGAGG	304	56	[35]
	ENTR	TGAACTACGTTTCGTTCTTCTGG			
<i>viaB</i>	ViaBF	CACGCACCATCATTTACCCG	738	57	[37]
	ViaBR	AACAGGCTGTAGCGATTTAGG			
DT 104	104F 104R	ATGCGTTTGGTCTCACAGCC GCTGAGGCCACGGATATTTA	102	56	[38]

<https://doi.org/10.1371/journal.pone.0274636.t001>

fragments for the primers and target genes [35,36] are listed in Table 1. Amplification products were visualized on a 2% agarose gel (Invitrogen®), stained with 5 mg/mL ethidium bromide, and visualized using a UV transilluminator (Major Science®; Saratoga, CA USA).

### Antibiogram and assessment of antimicrobial resistance and virulence genes

The antibiogram test was performed on all identified microorganisms as described by Kirby-Bauer [39] with a disk diffusion assay, using Mueller–Hinton agar (Acumedia®). The antibiotics tested were amoxicillin (10 µg), ampicillin (10 µg), nalidixic acid (30 µg), colistin (10 µg), cefazolin (30 µg), ceftazidime (30 µg), ciprofloxacin (5 µg), chloramphenicol (30 µg), doxycycline (30 µg), streptomycin (10 µg), gentamicin (10 µg), tetracycline (30 µg), and sulfonamide (30 µg). The results were based on the Clinical and Laboratory Standards Institute [40] halo parameters, except for colistin standards, for which used the parameters defined by the European Committee on Antimicrobial Susceptibility Testing [41]. The presence of 17 antimicrobial resistance genes were investigated using the oligonucleotide sequences described in Table 2.

For research on virulence genes, 12 virulence markers were selected based on their ability to cause lesions in the host organism [50,51]. The oligonucleotide annealing temperatures are listed in Table 3.

Amplification reactions were performed in a final volume of 25 µL, containing 2 U Taq DNA polymerase (Invitrogen®), 2 mM phosphated deoxyribonucleotides (Invitrogen®), 1× buffer (200 mM Tris-HCl, pH 8.4, 500 mM KCl; Invitrogen®), 3mM MgCl<sub>2</sub> (Invitrogen®), and 1 µM primers. The reaction was performed in a MyCycler thermal cycler (BioRad®) under the following conditions: initial denaturation at 94°C for 3 min; 30 cycles at 94°C for 30 sec, annealing temperature according to each primer for 30 sec, and 72°C for 30 sec, and a final cycle at 72°C for 10 min. Amplification products were visualized on a 2% agarose gel (Invitrogen®), stained with 5 mg/mL ethidium bromide, and visualized using a transilluminator (Major Science®).

### Evaluation of *in vitro* biofilm-forming capability

The *in vitro* biofilm-forming capability was evaluated as described by Agostinho Davanzo *et al.* [57] and Borges *et al.* [15].

The 96-well polystyrene titration microplates (Kartell®; Noviglio, Italy) containing the strains were incubated for 24 and 72 h at 37°C (near optimal temperature for target microorganism multiplication [58]), 24°C (average ideal temperature for extracellular polymeric matrix component expression [59,60]), and 10°C (maximum temperature recommended by

**Table 2. Resistance genes.** Oligonucleotides used for the antimicrobial resistance gene detection.

Antibiotic class	Gene	Primer	Nucleotide sequence (5'→3')	Size (bp)	Annealing temperature (°C)	Reference
Polymyxins	<i>MCR-1</i>	CLR5-F CLR5-R	CGGTCAGTCCGTTTGTTC CTTGGTCGGTCTGTAGGG	309	52	[42]
	<i>MCR-2</i>	MCR2-F MCR2-R	TGGTACAGCCCCTTATT GCTTGAGATTGGTTATGA	1617	48	[43]
	<i>MCR-3</i>	MCR3-F MCR3-R	TTGGCACTGTATTTGCATTT TTAACGAAATTGGCTGGAAACA	542	52	[44]
	<i>MCR-4</i>	Mcr-4 FW Mcr-4 RV	ATTGGGATAGTCGCCTTTTT TTACAGCCAGAATCATTATCA	487	51	[45]
Tetracyclines	<i>tet(A)</i>	tet(A)-F tet(A)-R	GTGAAACCCCAACATACCCC GAAGGCAAGCAGGATGTAG	887	53	[46]
	<i>tet(B)</i>	tet(B)-F tet(B)-R	CCTTATCATGCCAGTCTTGC ACTGCCGTTTTTTCGCC	773	53	[46]
	<i>tet(C)</i>	tet(C)-F tet(C)-R	ACTTGGAGCCACTATCGAC CTACAATCCATGCCAACCC	880	53	[46]
	<i>tet(M)</i>	tet(M)-1 tet(M)-2	GTTAAATAGTGTCTTGGAG CTAAGATATGGCTCTAACAA	700	49	[47]
Macrolides	<i>ermA</i>	ermA-F ermA-R	TCTAAAAAGCATGAAAAGAA CTTCGATAGTTTATTAATATTAGT	645	51	[48]
	<i>ermB</i>	ermB-F ermB-R	GAAAAGGTACTIONCAACCAATA AGTAACGGTACTTAAATGTTTAC	639	54	[48]
	<i>ermC</i>	ermC-F ermC-R	TCAAAACATAATATAGATAAA GCTAATATTGTTAAATCGTCAAT	642	51	[48]
	<i>ereA</i>	ere(A)-F ere(A)-R	GCCGGTGCTCATGAACCTGAG CGACTCTATTTCGATCAGAGGC	419	59	[46]
Amphenicols	<i>catI</i>	CATIF CATIR	AGTTGCTCAATGTACCTATAACC TTGTAATTCATTAAGCATTCTGCC	547	58	[46]
	<i>cmlA</i>	cmlA-F cmlA-R	CCGCCACGGTGTGTGTTATC CACCTTGCCTGCCCATCATTAG	698	58	[46]
Sulfonamide	<i>sulI</i>	sulI-F sulI-R	TTCGGCATTTCTGAATCTCAC ATGATCTAACCCCTCGGTCTC	822	53	[46]
β-lactams	<i>blaSHV</i>	blaSHV-F blaSHV-R	TCGCCTGTGTATTATCTCCC CGCAGATAAATCACCACAATG	768	51	[46]
	<i>ampC</i>	AmpC-For AmpC-Rev	TTCTATCAAMACTGGCARCC CCYTTTTATGTACCCAYGA	550	49	[49]
Aminoglycosides	<i>aac(3)-I</i>	aac(3)-I-F aac(3)-I-R	ACCTACTCCCAACATCAGCC ATATAGATCTCACTACGCGC	157	54	[46]

<https://doi.org/10.1371/journal.pone.0274636.t002>

the Brazilian Federal Inspection Service for facilities intended for the roasting and deboning of carcasses from cooling [61]).

The mean absorbance obtained from triplicate readings was used to determine the final optical density of each strain (ODf), which was compared with that of the negative control (ODn). The isolates were categorized into non-biofilm-forming isolates (NF) when  $ODf \leq ODn$ , weakly biofilm-forming when  $ODn < ODf \leq 2 \times ODn$ , moderate biofilm-forming when  $2 \times ODn < ODf \leq 4 \times ODn$ , or strong biofilm-forming when  $4 \times ODn < ODf$  [62].

Statistical analyses were performed using SAS software (v9.4; Cary, NC, USA) at 5% significance level. Initially, a normality test was performed (Shapiro–Wilk), and the data were subjected to analysis of variance using PROC GLIMMIX. The variables included time, detection points, temperature, and their interactions.

This study did not require permission from an ethics committee as no human or animal experimentation was involved.

**Table 3. Virulence genes and *E. coli* serotypes.** Oligonucleotides used for virulence gene and serotype detection in *E. coli* strains.

Gene	Primer	Oligonucleotides sequence (5'→3')	Size (bp)	Annealing temperature (°C)	References
<i>Tir α</i>	B139 B152	CAGCCTGCCACTTACCTTCACA CGCTAACCTCCAAACCATT	781	54.2	[52]
<i>Tir β</i>	B139 B140	CAGCCTGCCACTTACCTTCACA TGTATGTCGCACTCTGATT	342	53.4	[52]
<i>Tir γ</i>	B139 B141	CAGCCTGCCACTTACCTTCACA GTCGGCAGTTTCAGTTTCAC	560	54.7	[52]
<i>Stx1</i>	stx1F stx1R	AGAGCGATGTTACGGTTTG TTGCCCCAGAGTGGATG	388	50	[53]
<i>Stx2</i>	stx2F stx2R	TGGGTTTTTCTTCGGTATC GACATCTGGTTGACTCTCTT	807	45	[53]
<i>Eae</i>	eaeAF eaeAR	AGGCTTCGTCACAGTTG CCATCGTCACCAGAGGA	570	48	[53]
<i>hlyA</i>	hlyAF hlyAR	GCATCATCAAGCGTACGTTCC AATGAGCCAAGCTGGTTAAGCT	534	57	[54]
<i>Saa</i>	SAADF SAADR	CGTGATGAACAGGCTATTGC ATGGACATGCCTGTGGCAAC	119	55	[55]
<i>EspP</i>	esp-A esp-B	AAACAGCAGGCACTTGAACG GGAGTCGTCAGTCAGTAGAT	1830	56	[56]
O111	O111F O111R	TAGAGAAATTATCAAGTTAGTTCC ATAGTTATGAACATCTTGTGTTAGC	406	60	[54]
O113	O113F O113R	AGCGTTTCTGACATATGGAGTG GTGTTAGTATCAAAAGAGGCTCC	593	60	[56]
O157	O157F O157R	CGGACATCCATGTGATATGRG TTGCCATGTACAGCTAATCC	259	60	[54]

<https://doi.org/10.1371/journal.pone.0274636.t003>

## Results

### *E. coli* detection and isolation

Twenty-one (47.72%) *E. coli* strains were detected in the swabs collected from the environment, utensils, and equipment of swine slaughterhouses, with 9 and 12 isolates being obtained from slaughterhouses A and B, respectively, between 2019 and 2021. The detection points for each *E. coli* isolate, as well as the total number of isolates per collection point, are listed in

**Table 4.**

**Table 4. *E. coli* detection points.** Points of *E. coli* detection in the environment, equipment, and utensils of swine slaughterhouses A and B located in the Federal District of Brazil.

<i>E. coli</i> detection points in swine slaughterhouses	Visit 1 Slaughterhouse A	Visit 2 Slaughterhouse A	Visit 1 Slaughterhouse B	Visit 2 Slaughterhouse B	Total <i>E. coli</i> isolates
Chute of viscera	1	0	0	0	1
Drains (dirty area)	1	0	1	0	2
Bleeding knife	0	1	1	0	2
Drains (clean area)	0	1	1	0	2
Viscera table	1	0	1	0	2
Dehairing machine	1	1	0	0	2
Table	0	0	1	1	2
Carcass splitting saw	0	0	1	1	2
Clean area wall	0	0	1	0	1
Floor cooling chamber	1	0	1	1	3
Walls cooling chamber	0	1	1	0	2
Total swabs	5	4	9	3	21

<https://doi.org/10.1371/journal.pone.0274636.t004>



*E. coli* was most commonly detected from the swabs of the carcass cooling chamber floor (75% samplings), and least commonly from the swabs of the viscera kicker and the wall present in the clean area of the slaughter room (25% for both locations). *E. coli* was not detected in the toilet table, carcass splitting saw, or the clean area wall of the slaughter room of slaughterhouse A, as well as the viscera kicker and dehairing machine of slaughterhouse B during either visit.

### **Salmonella spp. detection and isolation**

Only one (2.27%) isolate of *Salmonella* spp. was detected from the 44 swab samples collected from swine slaughterhouses A and B. The isolate was recovered in the carcass cooling chamber during the second visit to slaughterhouse A.

The *Salmonella* genus was confirmed (204-bp fragment) and the *S. Typhi* serotype (738-bp fragment) was identified by colony PCR (Fig 1).

### **L. monocytogenes detection**

*L. monocytogenes* was not detected in any of the swab samples in the present study.


### **Antibiogram and resistance genes of E. coli isolates**

All 21 *E. coli* isolates were resistant or showed intermediate sensitivity to 12 of the 13 antimicrobials tested; 20 (95.2%) isolates were resistant to ampicillin and chloramphenicol each, 18 (85.8%) to amoxicillin, 17 (80.95%) to streptomycin and tetracycline each, 13 (61.9%) to sulfonamide, 12 (57.15%) to nalidixic acid and doxycycline each, 11 (52.4%) to cefazolin, seven (33.3%) to ciprofloxacin, five (23.8%) to gentamicin, and two (9.52%) to colistin. Moreover, six (28.6%) isolates presented intermediate resistance to ciprofloxacin, four (19.05%) to streptomycin, two (9.52%) to nalidixic acid, and one (4.8%) to chloramphenicol, cefazolin, and gentamicin each. None of the 21 isolates tested was resistant to ceftazidime (Table 5).

Twelve isolates expressed a resistance phenotype, and the antibiogram results were confirmed by a resistance gene detection (Table 6). Isolates 1 and 6 were resistant to ampicillin and tetracycline in the antibiogram and possess the respective genes *ampC* and *tet(A)*; isolates 10 and 17 were resistant to tetracycline in the antibiogram and possess the *tet(A)* gene; isolates 5 and 32 were resistant to chloramphenicol in the antibiogram and possess the genes *clmA* and *catI*; isolate 14 was resistant to tetracycline and sulfonamide in antibiogram and possess *tet(A)*, *tet(B)*, and *sull* genes; isolate 15 was resistant to tetracycline and colistin and possess *MCR-1*, *MCR-3*, and *tet(B)* genes; isolate 32 was resistant to chloramphenicol and possess the *catI* gene; isolate 33 was resistant to ampicillin and possess the genes *ampC* and *blaSHV*; isolate 41 was resistant to ampicillin and tetracycline and possess the genes *ampC* and *tet(B)*; and isolate 43 was also resistant to ampicillin and possess the gene *ampC*. The isolates 1, 10, and 40 possess the resistance genes *sull*, *MCR-3*, and *tet(B)*, respectively, but were sensitive to sulfonamide and colistin in the antibiogram. In this study, we could not relate the *aac(3)-I* to the aminoglycosides and *ermA*, *ermB*, *ermC*, and *ereA* to the macrolides.

### **Antibiogram and resistance genes of Salmonella spp.**

The sole *S. Typhi* isolate was resistant to 8 of the 13 antimicrobials tested, including nalidixic acid, cefazolin, chloramphenicol, doxycycline, streptomycin, gentamicin, tetracycline, and sulfonamide. The isolate was sensitive to amoxicillin, ampicillin, ciprofloxacin, ceftazidime, and colistin. Intermediate resistance to any of the investigated antimicrobials was not detected. Moreover, the antimicrobial resistance gene, *ampC*, which corresponds to  $\beta$ -lactams, as well as *tet(B)*, *tet(C)*, and *tet(M)*, which corresponds to tetracyclines, were detected. The resistance to



**Fig 1. *Salmonella* Typhi.** PCR confirmation of *S. Typhi* isolated from slaughterhouse A located in the Federal District of Brazil. 1) 100-bp marker (Invitrogen®), 2) negative control, 3) positive control for *Salmonella* spp., 204-bp fragment (*ompC* primer), 4) 204-bp fragment (*ompC* primer) for *Salmonella* spp. and 738-bp fragment (*viaB* primer) for Typhi serotype. Visualization on a 2% agarose gel stained with 0.5 µg/mL ethidium bromide in an ultraviolet transilluminator (Major Science®).

<https://doi.org/10.1371/journal.pone.0274636.g001>

tetracycline and doxycycline were confirmed by the presence of *tet(B)*, *tet(C)*, and *tet(M)*. The *Salmonella* spp. isolate was sensitive to ampicillin, an antibiotic of  $\beta$ -lactam class, and possesses the *ampC* gene, indicating  $\beta$ -lactam resistance (Table 7).

Additionally, *S. Typhi* presented an antimicrobial-resistant phenotype to cefazolin, nalidixic acid, chloramphenicol, sulfonamide, streptomycin, gentamicin, tetracycline, and doxycycline. However, no resistant genes were investigated in this study against the drugs, such as *cat1* and *clmA* for chloramphenicol (amphenicols), *aac(3)-I* for streptomycin and gentamicin (aminoglycosides), or *sul1* for sulfonamides.

In addition, no resistance genes for polymyxins (*MCR-1*, *MCR-2*, *MCR-3*, and *MCR-4*) or macrolides (*ermA*, *ermB*, *ermC*, and *ereA*) were detected.



**Table 5. *E. coli* antibiograms.** Antibiogram results of 21 *E. coli* isolates from swine slaughterhouses A and B.

Antibiotic class	Antimicrobial	Number of resistant isolates (%)	Number of intermediate resistance isolates (%)	Number of sensitive isolates (%)	Total resistant and intermediate isolates (%)
Polymyxins	Colistin (COL)	2 (9.52%)	0 (0%)	19 (90.48%)	2 (9.52%)
Tetracyclines	Tetracycline (TET)	17 (80.95%)	0 (0%)	4 (19.05%)	17 (80.95%)
	Doxycycline (DOX)	12 (57.15%)	2 (9.52%)	7 (33.33%)	14 (66.7%)
Amphenicols	Chloramphenicol (CLO)	20 (95.2%)	1 (4.8%)	0 (0%)	21 (100%)
Sulfonamides	Sulfonamide (SUL)	13 (61.9%)	0 (0%)	8 (38.1%)	13 (61.9%)
$\beta$ -lactams	Amoxicillin (AMO)	18 (85.8%)	0 (0%)	3 (14.2%)	18 (85.8%)
	Ampicillin (AMP)	20 (95.2%)	0 (0%)	1 (4.8%)	20 (100%)
Aminoglycosides	Streptomycin (EST)	17 (80.95%)	4 (19.05%)	0 (0%)	21 (100%)
	Gentamicin (GEN)	5 (23.8%)	1 (4.8%)	15 (71.4%)	6 (28.6%)
Cephalosporins	Cefazolin (CFZ)	11 (52.4%)	1 (4.8%)	9 (42.8%)	12 (57.15%)
	Ceftazidime (CAZ)	0 (0%)	0 (0%)	21 (100%)	0 (0%)
Fluoroquinolones	Ciprofloxacin (CIP)	7 (33.3%)	6 (28.6%)	8 (38.1%)	13 (61.9%)
	Nalidixic acid (NAL)	12 (57.15%)	2 (9.52%)	7 (33.33%)	14 (66.7%)

<https://doi.org/10.1371/journal.pone.0274636.t005>

### Virulence genes in *E. coli* isolates

Seven of the nine investigated virulence genes were detected in the *E. coli* isolates. Thirteen (61.9%) of the 21 *E. coli* isolates presented at least one virulence gene, of which five (23.8%), isolates harbored *tir*  $\alpha$ , five (23.8%) harbored *tir*  $\beta$ , five (23.8%) harbored *stx-1*, three (14.2%) harbored *tir*  $\gamma$ , three (14.2%) harbored *hlyA*, one (4.8%) harbored *ea*e, and one (4.8%) presented *saa*. Virulence genes *stx-2* and *Esp* were not detected.

As for the investigated serotypes, two (9.5%) isolates presented serotype *O157* (isolates 15 and 22). *O111* and *O113* serotypes were not detected in this study. The individual isolate results, detection point in the industry, antibiogram results, and detection results of antimicrobial resistance genes are presented in [Table 3](#).

### Evaluation of *in vitro* biofilm formation capacity of *E. coli* isolates

Biofilm-forming capacity increased after incubating for 72 h, and the optical density at 24 h indicated an initial stage of adherence. Biofilm-forming capacity was the highest after incubating at 10°C, while it was the lowest after incubating at 37°C for 24 h. After incubating for 72 h at 37°C, 2 (9.5%), 6 (28.6%), 11 (52.4%), and 2 (9.5%) *E. coli* isolates showed strong, moderate, weak, and no biofilm-forming capacity, respectively. Interestingly, 4 (19.05%), 10 (47.75%), 6 (28.6%), and 1 (4.8%) isolates showed strong, moderate, weak, and no biofilm-forming capacity, respectively, at 24°C; furthermore, 4 (19.05%), 8 (38.1%), 7 (33.3%), and 2 (9.5%) isolates showed strong, moderate, weak, and no biofilm-forming capacity, respectively, at 10°C. According to the statistical analyses performed, biofilm formation capacity was significantly different at the 5% significance level ( $P < 0.0001$ ) in relation to different temperatures, incubation periods, and swab detection points.

Individual identification, as well as the optical density and classification of biofilm-forming capacity of the 21 *E. coli* isolates after incubating at the three temperatures for 24 and 72 h are presented in [S1 Table](#). Concerning the detection points, isolate 40 had the highest biofilm-forming capacity, while isolate 22 (isolated from the carcass cooling chamber wall) did not form biofilms at any time or temperature conditions.

**Table 6. *E. coli* genes' detection and antibiograms.** Results of 21 *E. coli* antibiograms, detection of resistance and virulence genes, and detection points in slaughter-houses A and B.

<i>E. coli</i> isolate	Swine slaughterhouse	Detection point	Antibiogram Resistant	Antibiogram Intermediate resistance	Antibiogram Sensitive	Resistance genes	Virulence genes
1	A	Chute of visceras	NAL, AMO, AMP, CFZ, CLO, EST, TET	–	CAZ, CIP, COL, DOX, GEN, SUL	<i>ampC</i> , <i>tet(A)</i> , <i>sulI</i>	<i>hlyA</i> , <i>tir β</i> , <i>stx-1</i>
2	A	Drains (clean area)	NAL, AMO, AMP, CLO, DOX, EST, TET, SUL	CFZ	CAZ, CIP, COL, GEN	–	<i>eae</i> , <i>hlyA</i> , <i>tir α</i> , <i>tir β</i>
5	A	Evisceration table	AMO, AMP, CFZ, CLO, DOX, TET, SUL	EST, NAL	CAZ, CIP, COL, GEN	<i>clmA</i>	<i>Stx-1</i> , <i>tir-β</i>
6	A	Dehairing machine	NAL, AMO, AMP, CFZ, CLO, DOX, TET, SUL	CIP, EST	CAZ, COL, GEN	<i>ampC</i> , <i>tet(A)</i>	<i>tir α</i> , <i>tir γ</i>
10	A	Floor (cooling chamber)	NAL, AMO, AMP, CFZ, CIP, CLO DOX, EST, GEN, TET, SUL	–	CAZ, COL	<i>MCR-3</i> , <i>tet(A)</i>	<i>tir α</i>
14	A	Bleeding knife	NAL, AMO, AMP, CIP, CLO, DOX, EST, TET, SUL	–	CFZ, CAZ, COL, GEN	<i>tet(A)</i> , <i>tet(B)</i> , <i>sulI</i>	–
15	A	Drains (clean area)	NAL, AMO, AMP, CFZ, CIP, CLO, COL, DOX, EST, GEN, TET, SUL	–	CAZ	<i>MCR-1</i> , <i>MCR-3</i> , <i>tet(B)</i>	<i>tir α</i>
17	A	Dehairing machine	NAL, AMO, AMP, CFZ, CIP, CLO, DOX, EST, GEN, TET, SUL	–	CAZ, COL	<i>tet(A)</i>	<i>Saa</i>
22	A	Walls (cooling chamber)	AMO, AMP, CFZ, DOX, EST, TET	NAL, CIP, CLO	CAZ, COL, GEN, SUL	–	<i>tir α</i> , <i>tir β</i> , <i>tir γ</i>
24	B	Drains (dirty area)	AMO, AMP, CFZ, CIP, CLO, EST, TET, SUL	DOX	NAL, CAZ, COL, GEN	–	–
25	B	Bleeding knife	CLO, EST, TET	DOX	NAL, AMO, AMP, CAZ, CFZ, CIP, COL, GEN, SUL	–	<i>stx-1</i>
26	B	Drains (clean area)	NAL, AMO, AMP, CIP, CLO, DOX, EST, GEN, TET, SUL	–	CFZ, CAZ, COL	–	–
27	B	Evisceration table	NAL, AMO, AMP, CLO, COL, EST, TET, SUL	–	CFZ, CAZ, CIP, DOX, GEN	–	–
29	B	Table (dirty area)	AMO, AMP, CLO, EST, TET, SUL	–	NAL, CFZ, CAZ, CIP, COL, DOX, GEN	–	–
30	B	Carcass splitting saw	AMO, AMP, CLO, EST, TET	CIP	NAL, CFZ, CAZ, COL, DOX, GEN, SUL	–	–
31	B	Wall (dirty area)	NAL, AMO, AMP, CLO, DOX	CIP, EST	CFZ, CAZ, COL, GEN, TET, SUL	–	<i>Stx-1</i>
32	B	Floor (cooling chamber)	AMP, CFZ, CLO, EST	–	NAL, AMO, CAZ, CIP, COL, DOX, GEN, TET, SUL	<i>cat1</i>	<i>Stx-1</i>
33	B	Wall (cooling chamber)	AMP, CFZ, CLO	EST	NAL, AMO, CAZ, CIP, COL, DOX, GEN, TET, SUL	<i>ampC</i> , <i>blaSHV</i>	–
40	B	Table (dirty area)	NAL, AMO, AMP, CFZ, CLO, EST	CIP, GEN	CAZ, COL, DOX, TET, SUL	<i>tet(B)</i>	<i>tir γ</i>
41	B	Carcass splitting saw	NAL, AMO, AMP, CIP, CLO, DOX, EST, GEN, TET, SUL	–	CFZ, CAZ, COL	<i>ampC</i> , <i>tet(B)</i>	<i>tir β</i>
43	B	Floor (cooling chamber)	AMO, AMP, CLO, DOX, EST, TET	CIP	NAL, CFZ, CAZ, COL, GEN, SUL	<i>ampC</i>	–

\* The inhibition zone diameters were measured and interpreted according to the CLSI [40] parameters, except for the standards for colistin, in which the parameters were defined by EUCAST [41].

\*\* In isolates without resistance/virulence genes, consider only the ones detected by the primers used in this study.

<https://doi.org/10.1371/journal.pone.0274636.t006>

**Table 7. *Salmonella* spp. results of the antibiogram.** Results of the *Salmonella* spp. isolate antibiogram by disk diffusion, antimicrobial resistance gene detection of *Salmonella* spp. isolate, and point of isolation point at swine slaughterhouse A located in the Federal District of Brazil.

<i>Salmonella</i> Typhi	Swine slaughterhouse identification	Detection point	Antibiogram resistance	Antibiogram sensitivity	Resistance genes detected
21	A	Floor (cooling chamber)	CFZ		
			NAL	AMP	<i>ampC</i>
			CLO	AMO	-
			SUL	CIP	-
			EST	CAZ	-
			GEN	COL	-
			DOX	-	-
			TET	-	<i>tet(B)</i> , <i>tet(C)</i> , <i>tet(M)</i>

<https://doi.org/10.1371/journal.pone.0274636.t007>

### Evaluation of *in vitro* biofilm formation capacity of *Salmonella* spp. isolates

*S. Typhi* incubated for 24 h at 37, 24, and 10°C showed weak biofilm-forming capacity at 37 and 24°C and did not form biofilms at 10°C. Moderate biofilm formation was observed when incubated for 72 h at 37°C, and weak biofilm formation was observed at 24 and 10°C. The optical densities as well as the biofilm-forming capacities of *S. Typhi* are described in [Table 8](#).

### Discussion

#### *E. coli* isolation from swine slaughterhouses located in the Federal District of Brazil

The microorganisms detected from the dehairing machine, tables, carcass splitting saw, and carcass cooling chamber floor swabs collected during both visits at slaughterhouses A and B suggest *E. coli* permanence and distribution in the slaughter process, corroborating the presence of *E. coli* on floors, tables, and knives of swine slaughterhouses in Nigeria [63]. In addition, Namvar & Warriner [64] detected the permanence of *E. coli* on swine slaughterhouse floors in two swabs collected on different dates. In general, the presence of *E. coli* in the swine slaughterhouse environment may indicate cleaning process failure [65]. Moreover, repeated isolation from the same industrial collection points may suggest the presence of bacterial biofilms [64]. Even a one-time *E. coli* recovery may indicate cross-contamination [66]. The presence of *E. coli* in dehairing machines in this study may be due to the presence of microorganisms in pig bristles, which are directed to the dehairing machines after slaughter and can contaminate the water and blades of the equipment [65]. A failure in equipment sanitization procedures may also cause bacterial contamination.

**Table 8. *Salmonella* Typhi biofilm formation.** *In vitro* biofilm-forming capacity of *S. Typhi* after 24 and 72 h incubation at 37, 24, and 10°C.

<i>Salmonella</i> isolate	Incubation period	ODf at 37°C	ODf at 24°C	ODf at 10°C	Classification at 37°C	Classification at 24°C	Classification at 10°C
21	24h	0.089	0.112	0.084	weak	weak	NF
	72h	0.135	0.074	0.084	moderate	weak	weak

\* The classification is based on the parameters described by Stepanović *et al.* [62], where ODf is the final optical density of the isolates, and ODn is the negative control optical density. ODn = 0.064 and 0.086 in isolates incubated for 24 and 72 h, respectively. The isolates were classified into non-biofilm-forming (NF, ODf ≤ ODn), weak biofilm-forming (ODn < ODf ≤ 2× ODn), moderate biofilm-forming (2× ODn < ODf ≤ 4× ODn), or strong biofilm-forming (4× ODn < ODf) according to their biofilm-forming ability and intensity.

<https://doi.org/10.1371/journal.pone.0274636.t008>

## ***Salmonella* spp. isolation in swine slaughterhouses located in the Federal District of Brazil**

*Salmonella* spp. have also been detected in the environment, equipment, and utensils of swine slaughterhouses in other European countries, such as Italy [67], Belgium [68], and the Netherlands [69], with frequent contamination points being the carcass splitting saw and knives used. The presence of *Salmonella* spp. in the carcass cooling chamber may be related to cross-contamination during the slaughter process, swine carcass cooling, and failures related to hygiene procedures. Botteldoorn *et al.* [68] have discussed the difficulty in stating the origin of the contamination site, since it may vary depending on the number of animals slaughtered daily in the establishment, pig farming practices responsible for raising and breeding domestic pigs as livestock for slaughter, failures in the conduction of standard sanitation operating procedures, and even failures in employee training. *S. Typhi* detection in the Brazilian slaughterhouses in this study is relevant because of the possibility of carcass cross-contamination when stored in the cooling chamber. This goes against Normative Instruction 79 [70], which fosters the importance of this specific microbiological analysis when approving risk-based ante- and post-mortem pig inspection procedures.

## **Studies on *L. monocytogenes* in swine slaughterhouses located in the Federal District of Brazil**

The non-isolation of *L. monocytogenes* from a swine slaughter facility in this study diverges from that reported by Moreno *et al.* [71] and Sereno *et al.* [72] in Brazil, Lariviere-Gauthier *et al.* [73] in Canada, Autio *et al.* [74] in Finland, and Morganti *et al.* [75] in Italy. A possible hypothesis for the non-detection of this microorganism would be the correct performance of the standard sanitation operating procedures in slaughterhouses, which may have been favored by the average size of the participating industries in this study, which slaughtered 110 animals per day, allowing better control of daily hygiene procedures. The non-detection of *L. monocytogenes* may also have occurred because of the restricted number of samples collected due to the resistance of the local industries participating in this study. However, it is important to emphasize that the non-detection of *L. monocytogenes* does not ensure its absence in slaughterhouses in the Federal District of Brazil since the microorganism presents a cosmopolitan characteristic [76,77]. Moreover, it has been detected in bovine meat cuts and the environment of bovine slaughterhouses in the Federal District [78]. In addition, this microorganism has also been detected in minced beef and hot dog sausages commercialized in this region [79].

## **Antibiogram and antimicrobial resistance gene detection in *E. coli* isolates**

The existence of multidrug-resistant isolates, such as isolates 10, 15, and 17, which are resistant to nalidixic acid, amoxicillin, ampicillin, cefazolin, ciprofloxacin, chloramphenicol, doxycycline streptomycin, gentamicin, tetracycline, and sulfonamide, is a public health concern as they may suggest the indiscriminate use of antibiotics for treatment, disease prevention, and growth promotion [80]. This may cause the emergence of resistant bacteria in livestock animals, their spread in the environment, or residues in animal products consumed by the population [80,81].

In this study, 95.2% (20/21) *E. coli* isolates were chloramphenicol-resistant, which is important because it is a broad-spectrum antibiotic with prohibited use in Brazil since 2003 according to Normative Instruction 09 [82]. The detection of resistant strains can be explained by the maintenance of resistance genes through co-selection with other resistance and virulence genes, often linked to transmissible/mobile genetic elements [83]. However, further studies should be

conducted to verify the possible origins of this antimicrobial resistance because only two *E. coli* isolates, 5 and 32, with *cat1* and *clmA* genes respectively, were chloramphenicol-resistant.

Chloramphenicol resistance has also been reported years after the ban (in the 1980s) on its use in animal feed in the USA. Chloramphenicol resistance was detected in 53% *E. coli* strains from diarrheic pigs, along with *clmA* [84]. This persistence is explained by the location of the *cmlA* in the class 1 integrons, allowing transfer by conjugation as they are linked to other genes encoding resistance to antimicrobials currently allowed for use in animal feed [84], which may also explain the resistance found in this study. In Japan, Harada *et al.* [85] corroborated this information by showing that *clmA* and *cat1* are involved in co-resistance, contributing to chloramphenicol-resistant strain selection, allowing it to persist despite its ban in swine feed, which may also explain the presence of resistance found in this study.

Isolate 15 was colistin resistant and possessed the resistance genes *MCR-1* and *MCR-3*, indicating polymyxin resistance. The World Health Organization (WHO) has classified polymyxins as critically important and the highest-priority antimicrobials [86]; thus, this result is concerning for public health. Other studies have shown that the use of this antibiotic increased in serious infection treatment in humans, and the presence of *MCR* genes confer transmissible resistance and spread resistant microorganisms through the food chain [80,87,88].

Some *E. coli* isolates showed antibiogram sensitivity and possessed resistance genes to the same antibiotic; isolate 1 harbored *sul1* and sensitivity to sulfonamide; isolate 10 was sensitive to colistin and contained *MCR-3*, whereas isolate 40 was sensitive to tetracycline and doxycycline, and *tet(B)* was detected, which can be explained by the non-expression of genes present in its bacterial or plasmid DNA [89].

### Antibiogram and antimicrobial resistance gene detection in *Salmonella* spp. isolates

As detected in the *E. coli* isolates, the resistance of *Salmonella* spp. to chloramphenicol is relevant to public health, since it has been banned from use in animal feed since 2003 [82]. This is problematic since food can be contaminated by resistant pathogens and distributed over large geographical areas, increasing antimicrobial resistance in the population that consumes such products [90]. Wu *et al.* [91] have also reported that *Salmonella* isolates from the environment and carcasses of pig slaughterhouses in China were chloramphenicol-resistant. Botteldoorn *et al.* [92] also detected a chloramphenicol-resistant microorganism in Belgian pig slaughterhouse environments, utensils, and carcasses.

As described previously, ampicillin ( $\beta$ -lactam) sensitivity and *ampC* ( $\beta$ -lactam) detection can be explained by the lack of expression of the gene present in its bacterial or plasmid DNA [93]. Consequently, findings such as phenotypic resistance to an antibiotic (amphenicols, aminoglycosides, and sulfonamides), and non-detection of correlated resistant genes, cannot be interpreted as the absence of resistance genes. The presence of cross-resistance [94,95] is considered valid in this case. Another possibility is the inappropriate methodology of primer choice, since classic primers were used in gene detection, and others such as *sul2*, *sul3*, and *floR* [96], were not included in this study. Furthermore, Schwan *et al.* [97] and Jearnsripong *et al.* [98] showed a concordance of phenotypic and genotypic AMR results of *Salmonella* spp. that represented the results different from those of this study. Therefore, to elucidate the origin of phenotype resistance, complete genome sequencing would be required.

### Virulence genes in *E. coli* isolates

*E. coli* isolate 2 possessed the highest number of virulence genes, including *hlyA*, *eae*, *tir*  $\alpha$ , and *tir*  $\beta$ , which are associated with the antibiogram profile of resistance to nalidixic acid,

amoxicillin, ampicillin, chloramphenicol, doxycycline, streptomycin, tetracycline, and sulfonamide. These findings imply the importance of *E. coli* isolate 2 due to the public health risks caused by the presence of these genes. *hlyA* encodes alpha-hemolysin exotoxin and is related to clinical infections in humans, such as pyelonephritis and sepsis [99]. *eae* and *tir* can be related to enteropathogenic *E. coli* strains, since *eae* encodes the adhesion factor intimin and *tir* is an intimin receptor, allowing the attaching and effacing pathogenesis mechanism, causing lesions in the intestinal mucosa of humans and animals [100,101]. The presence of these virulence genes in addition to the resistance to the antimicrobials mentioned suggests a potential risk to the population. Moreover, *E. coli* isolate 17 presented *saa* association with *tet(A)* and antibiogram resistance profile to nalidixic acid, amoxicillin, ampicillin, cefazolin, ciprofloxacin, chloramphenicol, doxycycline, streptomycin, gentamicin, tetracycline, and sulfonamide. It implies a risk to public health since the isolate is resistant to multiple antibiotics and possesses *saa*, which may lead to clinical cases of severe diarrhea in humans [102].

Additionally, it is important to highlight that 5/21 *E. coli* isolates detected in post-sanitation locations of processing plants were non-O157:H7 Shiga toxin-producing *E. coli* strains (STECs). The presence of *stx-1* in non-O157 strains, as observed in isolates 1, 5, 25, 31, and 32, even though they are not from a serotype conventionally associated with pathologies (O157:H7), is associated with severe disease in humans [103,104]. The *E. coli* virulence genes detected in this study confer pathogenicity and are a potential risk to public health [105]; the isolates were isolated from the surfaces of equipment, utensils, and the environment of swine slaughterhouses, having direct and/or indirect contact with the food produced. This may cause direct contamination or cross-contamination of final products that will be consumed by the population of the Federal District area and other Brazilian states.

The virulence genes present in isolates from slaughterhouses A and B were different, which can be attributed to the different batches of animals received for slaughter and different sanitary management in livestock animal farms [106]. Although very few studies have verified virulence genes in *E. coli* detected in the environment, equipment, and utensils in swine slaughterhouses/carcasses, some studies have detected these virulence genes in pig carcasses [107–110], suggesting that the virulence genes investigated in this study are circulating in *E. coli* strains in pigs. In addition to the potential public health risk related to food contamination, it is important to emphasize the economic loss due to infection by pathogenic *E. coli* strains in pigs (particularly piglets) and feed conversion reduction due to diarrheal symptoms, which may contaminate other pigs on the farm and cause death due to severe dehydration or the development of syndromes related to pathogenic *E. coli* strains [111].

### ***In vitro* evaluation of biofilm-forming capacity of *E. coli* isolates**

The biofilm formation in most *E. coli* isolates was the maximum after incubating for 72 h at 10°C. This corroborates with the guidelines of the Brazilian Ministry of Agriculture, Livestock and Supply Ordinance No. 1304 [112] about the importance of daily cleaning in slaughterhouses after the activities and before starting the slaughter process, aiming for hygienic sanitary quality of produced food. It is relevant that the highest capacity to form biofilms occurred at refrigeration temperature (10°C); this condition resembles that of the climate-controlled deboning in Brazilian slaughterhouses [112]. Therefore, *E. coli* strains forming biofilms at this temperature can represent a contamination risk of the final food product.

Five (23.8%) *E. coli* isolates showed strong biofilm-forming capacities in at least one of the three temperatures tested, among which, isolates 32 and 40 harbored *cat1* and *tet(B)*, respectively. Moreover, 15 (71.4%) isolates showed moderate biofilm-forming capacity at all temperatures tested and antimicrobial resistance genes were detected in nine of the 15 isolates: isolate



1 (*ampC* and *tet(A)*, and *sul1*), isolate 5 (*clmA*), isolate 6 (*tet(A)* and *ampC*), isolate 10 (*tet(A)* and *MCR-3*), isolate 17 (*tet(A)*), isolate 33 (*ampC* and *blaSHV*), isolate 40 (*tet(B)*), isolate 41 (*tet(B)* and *ampC*), and isolate 43 (*ampC*). These results show the importance of *E. coli* isolates due to the risk posed to public health and the capacity to spread antimicrobial resistance in the environment [113]. In addition to the presence of resistance genes, these isolates harbored virulence genes *stx-1*, *saa*, *hlyA*, and *tir*, which are associated with serious disease development in humans, reinforcing the potential risk to consumers of the meat processed in meatpacking industries.

Bacterial biofilm formation is a serious problem in food industries [114] as it allows microorganisms to remain viable for months on surfaces after sanitization and hygiene procedures, becoming a recurrent point of contamination [115]. The recurring failure of sanitization processes causes bacterial attachment to abiotic surfaces; once established in the environment, removing the biofilm is challenging in food industries as a self-produced extracellular matrix enables the adhesion of other microorganisms and the colonization of several surfaces [116]. In industrial environments, complex multi-species communities permit bacterial cell attachment and detachment, enabling product cross-contamination and, in turn, product shelf-life reduction and disease transmission [117]. This is the first study to evaluate the biofilm formation capacity of *E. coli* in swine slaughterhouses in Brazil.

### ***In vitro* evaluation of *Salmonella* spp. biofilm-forming capacity**

*S. Typhi* presented a moderate biofilm-forming capacity at 37°C after incubation for 72 h, which may have occurred because of its ideal growth temperature [118]. Moreover, the isolate showed weak biofilm-forming capacity or did not form biofilms at other temperatures and incubation periods even though it was detected in the environment. Very few studies have evaluated the biofilm-forming capacity of *Salmonella* spp. in Brazilian poultry slaughterhouses; the results of this study were similar to those reported by Garcia *et al.* [119], which reported weak and moderate biofilm-forming capacity of *Salmonella* strains isolated from poultry carcasses and equipment used in poultry farms in São Paulo. Sereno *et al.* [120] reported similar results, detecting weak and moderate biofilm-forming *Salmonella* strains on frozen poultry carcasses in Paraná. It is important to emphasize that even as a non-biofilm former (10°C after 24 h incubation), *S. Typhi* is the agent of typhoid fever, a disease widely described and clinically characterized by high fever, headache, diarrhea, and abdominal pain after consuming contaminated food [121,122]. Thus, this pathogen poses public health risk because it presents multidrug resistance and resistance genes (*tet(B)*, *tet(C)*, *tet(M)*, and *ampC*) and can attach to surfaces.

### **Conclusion**

This is the first study to evaluate the biofilm-forming capacity of *Salmonella* spp. isolated from a swine slaughterhouse in Brazil. Furthermore, 21 *E. coli* isolates and one *S. Typhi* isolate were detected in the environment and equipment. The *E. coli* isolates were multidrug-resistant and harbored resistance and virulence genes. Moreover, 23.8% and 71.4% *E. coli* isolates presented strong and moderate biofilm-forming capacity, respectively. The *S. Typhi* isolate was multidrug-resistant and possessed a tetracycline resistance gene. Additionally, it presented moderate biofilm-forming capacity at 37°C after incubating for 72 h. The results of this study suggest a public health risk. The association of the above-stated pathogens with foodborne diseases has been extensively documented, and the decrease in foodborne disease occurrences is closely related to increased food quality through careful hygienic actions within the industries. Reducing or eliminating pathogenic microorganisms before bacterial biofilm formation to ensure

the hygienic and sanitary quality of the final product is a guaranteed way to avoid public health risks. Furthermore, it is important to emphasize the risk of spreading resistance genes in the environment. The presence of multiple antimicrobial resistance genes in the isolates in this study indicates the need for the rational use of these drugs to preserve their effectiveness for future use.

## Supporting information

**S1 Fig. Raw gel image of Fig 1 - PCR confirmation of *Salmonella* Typhi.** Row 1) 100-bp marker (Invitrogen®), row 2) negative control, row 3) positive control for *Salmonella* spp., 204-bp fragment (ompC primer), row 4) 204-bp fragment (ompC primer) for *Salmonella* spp., and 738-bp fragment (viaB primer) for Typhi serotype. Visualization on a 2% agarose gel stained with 0.5 µg/mL ethidium bromide in an ultraviolet transilluminator (Major Science®).  
(PDF)

**S1 Table. *E. coli* biofilm formation.** Biofilm-forming capacity in 21 *E. coli* isolates incubated for 24 h and 72 h at three different temperatures (37, 24 and 10°C). The classification is based on the parameters described by Stepanović et al. [62], where ODF is the final optical density of the isolates, and ODn is the negative control optical density. ODn = 0.064 and 0.086 in isolates incubated for 24 h and 0.086 in isolates incubated for 72 h, respectively. The isolates were classified into non-biofilm-forming (NF) when  $ODf \leq ODn$ , weak biofilm-forming ( $ODn < ODF \leq 2 \times ODn$ ), moderate biofilm-forming ( $2 \times ODn < ODF \leq 4 \times ODn$ ), or strong biofilm-forming ( $4 \times ODn < ODF$ ) according to their biofilm-forming ability and intensity.  
(PDF)

## Acknowledgments

We thank the staff at the Laboratory of Food Microbiology of the University of Brasília for providing aid when necessary.

Supplemental data (S1 Table) can be found in the Supporting Information section.

## Author Contributions

**Data curation:** Rebecca Lavarini dos Santos, Bruno Stéfano Lima Dallago.

**Investigation:** Rebecca Lavarini dos Santos, Bruno Stéfano Lima Dallago.

**Methodology:** Rebecca Lavarini dos Santos, Emilia Fernanda Agostinho Davanzo, Joana Marchesini Palma, Virgílio Hipólito de Lemos Castro, Hayanna Maria Boaventura da Costa, Simone Percmanis.

**Software:** Bruno Stéfano Lima Dallago.

**Supervision:** Ângela Patrícia Santana.

**Writing – original draft:** Rebecca Lavarini dos Santos.

## References

1. Brasil. Ministério da Saúde. Departamento de Vigilância das Doenças Transmissíveis. Apresentação sobre Surtos de Doenças Transmitidas por Alimentos no Brasil. 2010 [Cited 2021 October 8] Available from: [https://bvsm.sau.gov.br/bvs/publicacoes/manual\\_integrado\\_vigilancia\\_doencas\\_alimentos.pdf](https://bvsm.sau.gov.br/bvs/publicacoes/manual_integrado_vigilancia_doencas_alimentos.pdf).

2. Adley C, Ryan MP. Antimicrobial Food Packaging. Edition: 1st. Chapter: The Nature and Extent of Foodborne Disease. Elsevier, 2016.
3. World Health Organization (WHO). Salmonella (non-typhoidal). Fact sheets, 2018. [Cited 2021 June 3] Available from: [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal)).
4. Eng S, Pusparajah P, Mutalib NA, Ser H, Chan K, Lee L. Salmonella: A review on pathogenesis, epidemiology and antibiotic resistance. *Frontiers in Life Science*. 2015; 8(3):284–293.
5. Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Boletim Epidemiológico, v.51. 2020. [Cited 2022 June 1] Available from: <https://www.gov.br/saude/pt-br/assuntos/saude-de-a-a-z/d/dtha/arquivos/atualizacao-sobre-notificacao-de-surto-de-dtha-no-sinan-net.pdf>.
6. Yang SC, Lin CH, Aljufally IA. Current pathogenic *Escherichia coli* foodborne outbreak cases and therapy development. *Arch Microbiol*. 2017; 199(6):811–825. <https://doi.org/10.1007/s00203-017-1393-y> PMID: 28597303
7. Newell DG, Koopmans M, Verhoef L, Duizer E, Aidara-Kane A, Sprong H, et al. Food-borne diseases—The challenges of 20 years ago still persist while new ones continue to emerge. *Int. J. Food Microbiol*. 2010; 139:3–15.
8. Satpathy S, Sen SK, Pattanaik S, Raut S. Review on bacterial biofilm: An universal cause of contamination. *Biocatalysis and Agricultural Biotech*. 2016; 7:56–66.
9. Wang R. Biofilms and Meat Safety: A Mini-Review. *J Food Prot*. 2019; 82 (1): 120–127. <https://doi.org/10.4315/0362-028X.JFP-18-311> PMID: 30702946
10. Koo OK, Mertz AW, Akins EL, Sirsat SA, Neal JA, Morawicki R, et al. Analysis of microbial diversity on deli slicers using polymerase chain reaction and denaturing gradient gel electrophoresis technologies. *Lett. Appl. Microbiol*. 2013; 56:111–119. <https://doi.org/10.1111/lam.12021> PMID: 23121623
11. Wang H, Ye K, Wei X, Cao J, Xu X, Zhou G. Occurrence, antimicrobial resistance and biofilm formation of *Salmonella* isolates from a chicken slaughter plant in China. *Food Control*. 2013; 33(2):378–384.
12. Lakicevic B, Nastasijevic I. *Listeria monocytogenes* in retail establishments: Contamination routes and control strategies. *Food Rev. Int*. 2017; 33:247–269.
13. Todd ECD, Notermans S. Surveillance of listeriosis and its causative pathogen *Listeria monocytogenes*. *Food Control*. 2011; 22:1484–1490.
14. Food and Drugs Administration (FDA). Food Code, 2013. [Cited 2022 June 2] Available from: <https://www.fda.gov/media/87140/download>.
15. Borges KA, Furian TQ, Sousa SS, Menezes R, Tondo EC, Salles CTP, et al. Biofilm formation capacity of *Salmonella* serotypes at different temperature conditions. *Pesq. Vet. Bras*. 2018; 38(1):71–76.
16. Oliveira MMM, Brugnera DF, Piccoli RH. Biofilmes microbianos na indústria de alimentos: uma revisão. *Rev Inst Adolfo Lutz*. 2010; 2010; 69(3):277–284.
17. Matereke LT, Okoh AI. *Listeria monocytogenes* Virulence, Antimicrobial Resistance and Environmental Persistence: A Review. *Pathogens*. 2020; 9(7):528.
18. Alizade H, Hosseini Teshnizi S, Azad M, Shojae S, Gouklani H, Davoodian P, et al. An overview of diarrheagenic *Escherichia coli* in Iran: A systematic review and meta-analysis. *J Res Med Sci*. 2019; 24:23. [https://doi.org/10.4103/jrms.JRMS\\_256\\_18](https://doi.org/10.4103/jrms.JRMS_256_18) PMID: 31007693
19. Vasudevan R. Biofilms: microbial cities of scientific significance. *J Microbiol Exp*. 2014; 1(3):84–98.
20. Associação Brasileira de Proteína Animal (ABPA). Relatório anual. 2020 [cited 2021 October 10]. Available from: [https://abpa-br.org/wp-content/uploads/2020/05/abpa\\_relatorio\\_anual\\_2020\\_portugues\\_web.pdf](https://abpa-br.org/wp-content/uploads/2020/05/abpa_relatorio_anual_2020_portugues_web.pdf).
21. Souza JF, Souza ACF, Costa FN. Retrospective study of outbreaks of foodborne diseases in the Northeast and State of Maranhão, from 2007 to 2019. *Research, Society and Development*. 2021; 10(1):e36010111728.
22. Barros MA, Nero LA, Silva LC, d'Ovidio L, Monteiro FA, Tamanini R, et al. *Listeria monocytogenes*: Occurrence in beef and identification of the main contamination points in processing plants. *Meat Sci*. 2007; 76(4):591–596. <https://doi.org/10.1016/j.meatsci.2007.01.016> PMID: 22061233
23. Cabral CC, Panzenhagen PH., Delgado KF, Silva GRA, Rodrigues DP, Franco RM, et al. Contamination of carcasses and utensils in small swine slaughterhouses by *Salmonella* in the northwestern region of the State of Rio de Janeiro, Brazil. *Journal of Food Protection*, 2017; 80(7):1128–1132. <https://doi.org/10.4315/0362-028X.JFP-16-387> PMID: 28585863
24. Nicolau AI, Bolocan AS. Sampling the Processing Environment for *Listeria*. In: Jordan K, Fox EM, Wagner M. *Listeria monocytogenes: Methods and Protocols, Methods in Molecular Biology*. New York: Humana Press; 2014, pp. 3–14.

25. Pan Y, Breidt F Jr, Kathariou S. Resistance of *Listeria monocytogenes* biofilms to sanitizing agents in a simulated food processing environment. *Appl Environ Microbiol*. 2006; 72(12):7711–7717. <https://doi.org/10.1128/AEM.01065-06> PMID: 17012587
26. Vestby LK, Møretrø T, Langsrud S, Heir E, Nesse LL. Biofilm forming abilities of *Salmonella* are correlated with persistence in fish meal and feed factories. *BMC Vet Res*. 2009; 5:20. <https://doi.org/10.1186/1746-6148-5-20> PMID: 19473515
27. Sandrini CNM, Pereira MA, Brod CS, Carvalho JB, Aleixo JAG. *Escherichia coli* verotoxigênica: isolamento e prevalência em 60 propriedades de bovinos de leite da região de Pelotas, RS, Brasil. *Rev. Ciência Rural*. 2007; 37(1):175–182.
28. Feng P, Weagant SW, Grant MA, Burkhardt W. *Bacteriological Analytical Manual* 8th Edition, 4th chapter, 1998. Enumeration of *Escherichia coli* and the Coliform Bacteria. [Cited 2021 October 3] Available from: <https://www.fda.gov/food/laboratory-methods-food/bam-chapter-4-enumeration-escherichia-coli-and-coliform-bacteria>.
29. Garrity GB, Staley JT. *Bergey's Manual of Systematic Bacteriology: Volume Two*. New York: Springer, 2005.
30. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa n° 40. Aprovar os métodos analíticos, isolamento e identificação da *Salmonella* na carne bovina, avicultura e produtos derivados de ovos. 2005. [Cited 2021 October 7] Available from: [https://www.normasbrasil.com.br/norma/instrucao-normativa-40-2005\\_75792.html](https://www.normasbrasil.com.br/norma/instrucao-normativa-40-2005_75792.html)
31. Ryser ET, Donnelly CW. *Listeria*. In: Salfinger Y, Tortorello M Lou, editors. *Compendium of Methods for the Microbiological Examination of Foods*. 5th ed. American Public Health Association, 2015.
32. Brasil. Ministério da Saúde. Manual Técnico de Diagnóstico Laboratorial da *Salmonella* spp. 2011. [Cited 2021 September 9] Available from: [https://bvsms.saude.gov.br/bvs/publicacoes/manual\\_tecnico\\_diagnostico\\_laboratorial\\_salmonella\\_spp.pdf](https://bvsms.saude.gov.br/bvs/publicacoes/manual_tecnico_diagnostico_laboratorial_salmonella_spp.pdf).
33. International Organization for Standardization (ISO). ISO 6579/2002: Microbiology of food and animal feeding stuffs—Horizontal method for the detection of *Salmonella* spp. 2002 [Cited 2021 September 5] Available from: <https://www.iso.org/standard/29315.html>.
34. Freitas CG, Santana AP, da Silva PH, Gonçalves VS, Barros MA, Torres FA, et al. PCR multiplex for detection of *Salmonella* Enteritidis, Typhi and Typhimurium and occurrence in poultry meat. *Int J Food Microbiol*. 2010; 139:1–2.
35. Alvarez J, Sota M, Vivanco AB, Perales I, Cisterna R, Rementeria A, et al. Development of a multiplex PCR technique for detection and epidemiological typing of *Salmonella* in human clinical samples. *J Clin Microbiol*. 2004; 42(4):1734–1738. <https://doi.org/10.1128/JCM.42.4.1734-1738.2004> PMID: 15071035
36. Kwang J, Littledike ET, Keen JE. Use of the polymerase chain reaction for *Salmonella* detection. *Lett Appl Microbiol*. 1996; 22(1):46–51. <https://doi.org/10.1111/j.1472-765x.1996.tb01106.x> PMID: 8588887
37. Kumar S, Balakrishna K, Batra HV. Detection of *Salmonella enterica* serovar Typhi (S. Typhi) by selective amplification of *invA*, *viaB*, *fliC-d* and *prt* genes by polymerase chain reaction in multiplex format. *Lett Appl Microbiol*. 2006; 42(2):149–154. <https://doi.org/10.1111/j.1472-765x.2005.01813.x> PMID: 16441380
38. Pritchett LC, Konkel ME, Gay JM, Besser TE. Identification of DT104 and U302 phage types among *Salmonella enterica* serotype Typhimurium isolates by PCR. *J Clin Microbiol*. 2000; 38(9):3484–3488. <https://doi.org/10.1128/JCM.38.9.3484-3488.2000> PMID: 10970411
39. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol*. 1966; 45(4):493–496. PMID: 5325707
40. Clinical and Laboratory Standards Institute (CLSI). CLSI M100—ED30:2020 Performance Standards for Antimicrobial Susceptibility Testing. 30 ed. 2020. [Cited 2021 October 15] Available from: [https://clsi.org/media/3481/m100ed30\\_sample.pdf](https://clsi.org/media/3481/m100ed30_sample.pdf).
41. European Committee on Antimicrobial Susceptibility Testing (Eucast). Version 11.0. 2021. [Cited 2021 October 12] Available from: <https://eucast.org>.
42. Liu YY Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016; 16(2):161–168. [https://doi.org/10.1016/S1473-3099\(15\)00424-7](https://doi.org/10.1016/S1473-3099(15)00424-7) PMID: 26603172
43. Xavier BB, Lammens C, Ruhai R, Kumar-Singh S, Butaye P, Goossens H, et al. Identification of a novel plasmid-mediated colistin-resistance gene, *mcr-2*, in *Escherichia coli*. *Euro Surveill*. 2016; 21(27): <https://doi.org/10.2807/1560-7917.ES.2016.21.27.30280> PMID: 27416987

44. Yin W, Li H, Shen Y, Liu Z, Wang S, Shen Z, et al. Novel Plasmid-Mediated Colistin Resistance Gene *mcr-3* in *Escherichia coli*. *mBio*. 2017; 8:e00543–17. <https://doi.org/10.1128/mBio.00543-17> PMID: 28655818
45. Carattoli A, Villa L, Feudi C, Curcio L, Orsini S, Luppi A, et al. Novel plasmid-mediated colistin resistance *mcr-4* gene in *Salmonella* and *Escherichia coli*, Italy 2013, Spain and Belgium, 2015 to 2016. *Euro Surveill*. 2017; 22(31):30589. <https://doi.org/10.2807/1560-7917.ES.2017.22.31.30589> PMID: 28797329
46. Van TT, Chin J, Chapman T, Tran LT, Coloe PJ. Safety of raw meat and shellfish in Vietnam: an analysis of *Escherichia coli* isolations for antibiotic resistance and virulence genes. *Int J Food Microbiol*. 2008; 124(3):217–223. <https://doi.org/10.1016/j.ijfoodmicro.2008.03.029> PMID: 18457892
47. Aarestrup FM, Agerso Y, Ahrens P, Jorgensen JCO, Madsen M, Jensen LB. Antimicrobial susceptibility and presence of resistance genes in *Staphylococci* from poultry. *Veterinary Microb*. 2000; 74(4):353–364. [https://doi.org/10.1016/s0378-1135\(00\)00197-8](https://doi.org/10.1016/s0378-1135(00)00197-8) PMID: 10831857
48. Sutcliffe J, Grebe T, Tait-Kamradt A, Wondrack L. Detection of erythromycin-resistant determinants by PCR. *Antimicrobial agents and chemotherapy*, 1996; 40(11):2562–2566. <https://doi.org/10.1128/AAC.40.11.2562> PMID: 8913465
49. Schwartz T, Kohnen W, Jansen B, Obst U. Detection of antibiotic-resistant bacteria and their resistance genes in wastewater, surface water, and drinking water biofilms. *FEMS Microbiol Ecol*. 2003; 43(3):325–335. <https://doi.org/10.1111/j.1574-6941.2003.tb01073.x> PMID: 19719664
50. Caprioli A, Morabito S, Brugère H, Oswald E. Enterohaemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. *Vet Res*. 2005; 36(3):289–311. <https://doi.org/10.1051/vetres:2005002> PMID: 15845227
51. Kaper J, Nataro J, Mobley H. Pathogenic *Escherichia coli*. *Nat Rev Microbiol*. 2004; 123–140. <https://doi.org/10.1038/nrmicro818> PMID: 15040260
52. China B, Goffaux F, Pirson V, Mainil J. Comparison of *eae*, *tir*, *espA* and *espB* genes of bovine and human attaching and effacing *Escherichia coli* by multiplex polymerase chain reaction. *FEMS Microb*. 1999; 178(1):177–182. <https://doi.org/10.1111/j.1574-6968.1999.tb13775.x> PMID: 10483737
53. China B, Pirson V, Mainil J. Typing of bovine attaching and effacing *Escherichia coli* by multiplex in vitro amplification of virulence-associated genes. *Appl Environ Microbiol*. 1996; 62(9):3462–5. <https://doi.org/10.1128/aem.62.9.3462-3465.1996> PMID: 8795238
54. Paton AW, Paton JC. Detection and characterization of Shiga toxin-producing *Escherichia coli* by using multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli* *hlyA*, *rfbO111*, and *rfbO157*. *J Clin Microbiol*. 1998; 36(2):598–602. <https://doi.org/10.1128/JCM.36.2.598-602.1998> PMID: 9466788
55. Paton AW, Woodrow MC, Doyle RM, Lanser JA, Paton JC. Molecular characterization of a Shiga toxin-producing *Escherichia coli* O113:H21 strain lacking *eae* responsible for a cluster of cases of hemolytic-uremic syndrome. *J Clin Microbiol*. 1999; 37(10):3357–3361. <https://doi.org/10.1128/JCM.37.10.3357-3361.1999> PMID: 10488206
56. Paton AW, Paton JC. Direct detection and characterization of Shiga toxin-producing *Escherichia coli* by multiplex PCR for *stx1*, *stx2*, *eae*, *ehxA*, and *saa*. *J Clin Microbiol*. 2002; 40(1):271–274. <https://doi.org/10.1128/JCM.40.1.271-274.2002> PMID: 11773130
57. Agostinho Davanzo EF, dos Santos RL, Castro VL, Palma JM, Pribul BR, Dallago BSL, et al. Molecular characterization of *Salmonella* spp. and *Listeria monocytogenes* strains from biofilms in cattle and poultry slaughterhouses located in the federal District and State of Goiás, Brazil. *PLoS ONE* 2021; 16(11): e0259687. <https://doi.org/10.1371/journal.pone.0259687> PMID: 34767604
58. Kim C, Wilkins K, Bowers M, Wynn C, Ndegwa E. Influence of pH and temperature on growth characteristics of leading foodborne pathogens in a laboratory medium and select food beverages. *Austin Food Sci*. 2018; 3(1):1031.
59. Hufnagel DA, Depas WH, Chapman MR. The Biology of the *Escherichia coli* Extracellular Matrix. *Microbiol Spectr*. 2014; 3(3): <https://doi.org/10.1128/microbiolspec.MB-0014-2014> PMID: 26185090
60. Combrouse T, Sadvovskaya I, Faille C, Kol O, Guérardel Y, Midelet-Bourdin G. Quantification of the extracellular matrix of the *Listeria monocytogenes* biofilms of different phylogenetic lineages with optimization of culture conditions. *J Appl Microbiol*. 2013; 114(4):1120–1131. <https://doi.org/10.1111/jam.12127> PMID: 23317349
61. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Portaria MAPA n° 711. Normas técnicas de instalações e equipamentos para abate e industrialização de suínos. Brasília, 1995. [Cited 2021 September 29] Available from: <https://www.defesa.agricultura.sp.gov.br/legislacoes/portaria-mapa-711-de-01-11-1995,755.html>.
62. Stepanović S, Vuković D, Dakić I, Savić B, Svabić-Vlahović M. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. *J Microbiol Methods*. 2000; 40(2):175–179. [https://doi.org/10.1016/s0167-7012\(00\)00122-6](https://doi.org/10.1016/s0167-7012(00)00122-6) PMID: 10699673



63. Egbule OS, Iweriebor BC, Odum E. Beta-Lactamase-Producing *Escherichia coli* isolates recovered from pig handlers in retail shops and abattoirs in selected localities in southern Nigeria: Implications for public health. *Antib.* 2020; 10(1):9. <https://doi.org/10.3390/antibiotics10010009> PMID: 33374204
64. Namvar A, Warriner K. Application of enterobacterial repetitive intergenic consensus-polymerase chain reaction to trace the fate of generic *Escherichia coli* within a high capacity pork slaughter line. *Int J Food Microbiol.* 2006; 108(2):155–163. <https://doi.org/10.1016/j.ijfoodmicro.2005.11.006> PMID: 16386814
65. Rivas T, Vizcaíno JA, Herrera FJ. Microbial contamination of carcasses and equipment from an Iberian pig slaughterhouse. *J Food Prot.* 2000; 63(12):1670–1675. <https://doi.org/10.4315/0362-028x-63.12.1670> PMID: 11131889
66. Santos RP, Ferreira LC. Avaliação microbiológica do ambiente, utensílios, superfícies e das mãos dos manipuladores em uma unidade de abate de suínos na cidade de Januária-MG. *Caderno De Ciências Agr.* 2017; 9(1):44–48.
67. Piras F, Fois F, Mazza R, Putzolu M, Delogu ML, Lochi PG, et al. *Salmonella* Prevalence and Microbiological Contamination of Pig Carcasses and Slaughterhouse Environment. *Ital J Food Saf.* 2014; 3(4): 4581. <https://doi.org/10.4081/ijfs.2014.4581> PMID: 27800371
68. Botteldoorn N, Heyndrickx M, Rijpens N, Grijspeerd K, Herman L. *Salmonella* on pig carcasses: positive pigs and cross contamination in the slaughterhouse. *J Appl Microbiol.* 2003; 95(5):891–903. <https://doi.org/10.1046/j.1365-2672.2003.02042.x> PMID: 14633017
69. Swanenburg M, Urlings HA, Snijders JM, Keuzenkamp DA, van Knapen F. *Salmonella* in slaughter pigs: prevalence, serotypes and critical control points during slaughter in two slaughterhouses. *Int J Food Microbiol.* 2001; 70(3):243–54. [https://doi.org/10.1016/s0168-1605\(01\)00545-1](https://doi.org/10.1016/s0168-1605(01)00545-1) PMID: 11764190
70. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa n° 79. Procedimentos de inspeção ante e post mortem de suínos com base em risco. 2018. [Cited 2021 September 20] Available from: [https://www.in.gov.br/materia/-/asset\\_publisher/Kujrw0TZC2Mb/content/id/55444279/do1-2018-12-17-instrucao-normativa-n-79-de-14-de-dezembro-de-2018-55444116](https://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/55444279/do1-2018-12-17-instrucao-normativa-n-79-de-14-de-dezembro-de-2018-55444116) [Accessed September 20, 2021].
71. Moreno LZ, Paixão R, de Gobbi DD, Raimundo DC, Porfida Ferreira TS, Micke Moreno A, et al. Phenotypic and genotypic characterization of atypical *Listeria monocytogenes* and *Listeria innocua* isolated from swine slaughterhouses and meat markets. *Biomed Res Int.* 2014; 2014:742032. <https://doi.org/10.1155/2014/742032> PMID: 24987702
72. Sereno MJ, Viana C, Pegoraro K, da Silva DAL, Yamatogi RS, Nero LA, et al. Distribution, adhesion, virulence and antibiotic resistance of persistent *Listeria monocytogenes* in a pig slaughterhouse in Brazil. *Food Microbiol.* 2019; 84:103234. <https://doi.org/10.1016/j.fm.2019.05.018> PMID: 31421784
73. Larivière-Gauthier G, Letellier A, Kérouanton A, Bekal S, Quessy S, Fournaise S, et al. Analysis of *Listeria monocytogenes* strain distribution in a pork slaughter and cutting plant in the province of Quebec. *Food Prot.* 2014; 77(12):2121–2128.
74. Autio T, Säteri T, Fredriksson-Ahomaa M, Rahkio M, Lundén J, Korkeala H. *Listeria monocytogenes* contamination pattern in pig slaughterhouses. *J Food Prot.* 2000; 63(10):1438–1442. <https://doi.org/10.4315/0362-028x-63.10.1438> PMID: 11041148
75. Morganti M, Scaltriti E, Cozzolino P, Bolzoni L, Casadei G, Pierantoni M, et al. Processing-dependent and clonal contamination patterns of *Listeria monocytogenes* in the cured ham food chain revealed by genetic analysis. *Appl Environ Microbiol.* 2015; 82(3):822–831. <https://doi.org/10.1128/AEM.03103-15> PMID: 26590278
76. Raheem D. Outbreaks of listeriosis associated with deli meats and cheese: an overview. *AIMS Microbiol.* 2016; 2(3):230–250.
77. Vallim DC, Barroso Hofer C, Lisboa R, Barbosa AV, Alves Rusak L, dos Reis CM, et al. Twenty Years of *Listeria* in Brazil: Occurrence of *Listeria* Species and *Listeria monocytogenes* serovars in food samples in Brazil between 1990 and 2012. *Biomed Res Int.* 2015; 2015:540204. <https://doi.org/10.1155/2015/540204> PMID: 26539507
78. Palma JM, Lisboa RC, Rodrigues DP, Santos AFM, Hofer E, Santana AP. Caracterização molecular de *Listeria monocytogenes* oriundas de cortes cárneos bovinos e de abatedouros frigoríficos de bovinos localizados no Distrito Federal, Brasil. *Pesq. Vet. Bras.* 2016; 36(10):957–964.
79. Andrade RR, Silva PHC, Souza NR, Murata LM, Gonçalves VSP, Santana AP. Ocorrência e diferenciação de espécies de *Listeria* spp. em salsichas tipo hot dog a granel e em amostras de carne moída bovina comercializadas no Distrito Federal. *Rev. Ciência Rural.* 2014; 44(1): 147–151.
80. Barlaam A, Parisi A, Spinelli E, Caruso M, Taranto PD, Normanno G. Global Emergence of Colistin-Resistant *Escherichia coli* in Food Chains and Associated Food Safety Implications: A Review. *J Food Prot.* 2019; 82(8):1440–1448. <https://doi.org/10.4315/0362-028X.JFP-19-116> PMID: 31339371



81. Pacheco-Silva E, Souza JR, Caldas ED. Resíduos de medicamentos veterinários em leite e ovos. *Quím. Nova*. 2014; 37(1):111–122.
82. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa n° 9. Proíbe a fabricação, a manipulação, o fracionamento, a comercialização, a importação e o uso dos princípios ativos: cloranfenicol e nitrofuranos. 2003. [Cited 2021 October 15] Available from: <https://www.diariodasleis.com.br/legislacao/federal/25330-proibe-a-fabricacao-a-manipulacao-o-fracionamento-a-comercializacao-a-importacao-e-o-uso-dos-principios-ativos-cloranfenicol-e-nitrofuranos-e-os-produtos-que-contenham-estes-principios-ativos-pa.html>.
83. Rosengren LB, Waldner CL, Reid-Smith RJ. Associations between antimicrobial resistance phenotypes, antimicrobial resistance genes, and virulence genes of fecal *Escherichia coli* isolates from healthy grow-finish pigs. *Appl Environ Microbiol*. 2009; 75(5):1373–1380. <https://doi.org/10.1128/AEM.01253-08> PMID: 19139228
84. Bischoff KM, White DG, Hume ME, Poole TL, Nisbet DJ. The chloramphenicol resistance gene *cmIA* is disseminated on transferable plasmids that confer multiple-drug resistance in swine *Escherichia coli*. *FEMS Microbiol Lett*. 2005; 243(1):285–291. <https://doi.org/10.1016/j.femsle.2004.12.017> PMID: 15668031
85. Harada K, Asai T, Kojima A, Ishihara K, Takahashi T. Role of coresistance in the development of resistance to chloramphenicol in *Escherichia coli* isolated from sick cattle and pigs. *Am J Vet Res*. 2006; 67(2):230–235. <https://doi.org/10.2460/ajvr.67.2.230> PMID: 16454626
86. World Health Organization (WHO). Critically important antimicrobials for human medicine, Ranking of medically important antimicrobials for risk management of antimicrobial resistance due to non-human use - 6th rev. 2018. [Cited 2021 October 15] Available from: <https://apps.who.int/iris/bitstream/handle/10665/312266/9789241515528-eng.pdf>.
87. Berglund B. Acquired Resistance to Colistin via Chromosomal And Plasmid-Mediated Mechanisms in *Klebsiella pneumoniae*. *Infectious Microbes & Diseases*. 2019; 1(1):10–19.
88. Li Z, Cao Y, Yi L, Liu JH, Yang Q. Emergent polymyxin resistance: End of an Era? *Open Forum. Infect Dis*. 2019; 6(10):ofz368. <https://doi.org/10.1093/ofid/ofz368> PMID: 31420655
89. Amer MM, Mekky HM, Amer AM, Fedawy HS. Antimicrobial resistance genes in pathogenic *Escherichia coli* isolated from diseased broiler chickens in Egypt and their relationship with the phenotypic resistance characteristics. *Vet World*. 2018; 11(8):1082–1088. <https://doi.org/10.14202/vetworld.2018.1082-1088> PMID: 30250367
90. Dowling A, O'Dwyer J, Adley CC. Alternatives to antibiotics: future trends. In: *Microbial pathogens and strategies for combating them: science, technology and education*. Mendez-Vilas, Ed. Espanha: Formatex Research Center; 2013 pp. 216–226.
91. Wu B, Ed-Dra A, Pan H, Dong C, Jia C, Yue M. Genomic investigation of *Salmonella* isolates recovered from a pig slaughtering process in Hangzhou, China. *Front Microbiol*. 2021; 12:704636. <https://doi.org/10.3389/fmicb.2021.704636> PMID: 34305874
92. Botteldoorn N, Herman L, Rijpens N, Heyndrickx M. Phenotypic and molecular typing of *Salmonella* strains reveals different contamination sources in two commercial pig slaughterhouses. *Appl Environ Microbiol*. 2004; 70(9):5305–14. <https://doi.org/10.1128/AEM.70.9.5305-5314.2004> PMID: 15345414
93. McMillan EA, Gupta SK, Williams LE, Jové T, Hiott LM, Woodley TA, et al. Antimicrobial resistance genes, cassettes, and plasmids present in *Salmonella enterica* associated with United States Food animals. *Frontiers in microb*. 2019; 10:832.
94. Périchon B, Courvalin P, Stratton CW. Antibiotic Resistance. *Encyclopedia of Microbiology*, Academic Press; 2019. pp. 127–139.
95. López HS, Olvera LG. Problemática del uso de enrofloxacin en la avicultura en México. *Veterinaria México*. 2000; 31(2):137–145.
96. Lopes GV, Michael GB, Cardoso M, Schwarz S. Antimicrobial resistance and class 1 integron-associated gene cassettes in *Salmonella enterica* serovar Typhimurium isolated from pigs at slaughter and abattoir environment. *Vet Microbiol*. 2016; 194:84–92. <https://doi.org/10.1016/j.vetmic.2016.04.020> PMID: 27142182
97. Schwan CL, Lomonaco S, Bastos LM, Cook PW, Maher J, Trinetta V, et al. Genotypic and phenotypic characterization of antimicrobial resistance profiles in non-typhoidal *Salmonella enterica* strains isolated from Cambodian informal markets. *Front. Microbiol*. 2021; 12:711472. <https://doi.org/10.3389/fmicb.2021.711472> PMID: 34603240
98. Jeamsripong S.; Li X.; Aly S.S.; Su Z.; Pereira R.V.; Atwill E.R. Antibiotic resistance genes and associated phenotypes in *Escherichia coli* and *Enterococcus* from cattle at different production stages on a dairy farm in central California. *Antibiotics*. 2021; 10:1042. <https://doi.org/10.3390/antibiotics10091042> PMID: 34572624

99. May AY, Gleason TG, Sawyer RG, Pruett TL. Contribution of *Escherichia coli* alpha-hemolysin to bacterial virulence and to intraperitoneal alterations in peritonitis. *Infect Immun*. 2000; 68(1):176–183. <https://doi.org/10.1128/IAI.68.1.176-183.2000> PMID: 10603385
100. Souza CO, Melo TRB, Melo CSB, Menezes EM, Carvalho AC, Monteiro LCR. *Escherichia coli* enteropatógena: uma categoria diarreogénica versátil. *Rev Pan-Amaz Saude*. 2016; 7(2):79–91.
101. Fröhlicher E, Krause G, Zweifel C, Beutin L, Stephan R. Characterization of attaching and effacing *Escherichia coli* (AEEC) isolated from pigs and sheep. *BMC Microbiol*. 2008; 8:144. <https://doi.org/10.1186/1471-2180-8-144> PMID: 18786265
102. Paton AW, Srimanote P, Woodrow MC, Paton JC. Characterization of Saa, a novel autoagglutinating adhesin produced by locus of enterochia effacement-negative Shiga-toxigenic *Escherichia coli* strains that are virulent for humans. *Infect Immun*. 2001; 69(11):6999–7009. <https://doi.org/10.1128/IAI.69.11.6999-7009.2001> PMID: 11598075
103. Menrath A, Wieler LH, Heidemanns K, Semmler T, Fruth A, Kemper N. Shiga toxin producing *Escherichia coli*: identification of non-O157:H7-Super-Shedding cows and related risk factors. *Gut Pathog*. 2010; 2(1):7. <https://doi.org/10.1186/1757-4749-2-7> PMID: 20618953
104. Bettelheim KA. The non-O157 shiga-toxigenic (verocytotoxigenic) *Escherichia coli*; under-rated pathogens. *Crit Rev Microbiol*. 2007; 33(1):67–87. <https://doi.org/10.1080/10408410601172172> PMID: 17453930
105. Beutin L, Fach P. Detection of Shiga Toxin-Producing *Escherichia coli* from Nonhuman Sources and Strain Typing. *Microbiol Spectr*. 2014; 2(3): <https://doi.org/10.1128/microbiolspec.EHEC-0001-2013> PMID: 26103970
106. Borges CA, Beraldo LG, Maluta RP, Cardozo MV, Guth BE, Rigobelo EC, et al. Shiga toxigenic and atypical enteropathogenic *Escherichia coli* in the feces and carcasses of slaughtered pigs. *Foodborne Pathog Dis*. 2012; 9(12):1119–25. <https://doi.org/10.1089/fpd.2012.1206> PMID: 23186549
107. Botteldoorn N, Heyndrickx M, Rijpens N, Herman L. Detection and characterization of verotoxigenic *Escherichia coli* by a VTEC/EHEC multiplex PCR in porcine feces and pig carcass swabs. *Res Microbiol*. 2003; 154(2):97–104.
108. Bouvet J, Montet MP, Rossel R, Le Roux A, Bavai C, Ray-Gueniot S, et al. Effects of slaughter processes on pig carcass contamination by verotoxin-producing *Escherichia coli* and *E. coli* O157:H7. *Int J Food Microbiol*. 2002; 77(1–2):99–108. [https://doi.org/10.1016/s0168-1605\(02\)00053-3](https://doi.org/10.1016/s0168-1605(02)00053-3) PMID: 12076043
109. Essendoubi S, Yang X, King R, Keenlside J, Bahamon J, Diegel J, et al. Prevalence and characterization of *Escherichia coli* O157:H7 on pork carcasses and in swine colon contents from provincially licensed abattoirs in Alberta, Canada. *J Food Prot*. 2020; 083(11):1909–1917. <https://doi.org/10.4315/JFP-20-146> PMID: 32584991
110. Martins RP, da Silva MC, Dutra V, Nakazato L, Leite DS. Preliminary virulence genotyping and phylogeny of *Escherichia coli* from the gut of pigs at slaughtering stage in Brazil. *Meat Sci*. 2013; 93(3):437–440.
111. Camargo LRP, Suffredini IB. Impacto causado por *Escherichia coli* na produção de animais de corte no Brasil: revisão de literatura. *J Health Sci*. 2015; 33(2): 193–197.
112. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Portaria MAPA n° 1304. Normas técnicas de instalações e equipamentos para abate e industrialização de suínos. Brasília, 2018. [Cited 2021 September 29] Available from: <https://www.defesa.agricultura.sp.gov.br/legislacoes/portaria-mapa-n-1304-de-7-de-agosto-de-2018,1172.html>.
113. Barilli E, Vismarra A, Villa Z, Bonilauri P, Bacci C. ESBL *E. coli* isolated in pig's chain: Genetic analysis associated to the phenotype and biofilm synthesis evaluation. *Int J Food Microbiol*. 2019; 289:162–167. <https://doi.org/10.1016/j.ijfoodmicro.2018.09.012> PMID: 30245289
114. Stocco CW, Almeida L, Barreto EH, Bittencourt JVM. Microbiological quality control in beef cattle processing. *Rev Esp*. 2017; 38:22.
115. Galié S, García-Gutiérrez C, Miguélez EM, Villar CJ, Lombó F. Biofilms in the Food Industry: Health aspects and control methods. *Frontiers in microb*. 2018; 9:898. <https://doi.org/10.3389/fmicb.2018.00898> PMID: 29867809
116. Karimi A, Karig D, Kumar A, Ardekani AM. Interplay of physical mechanisms and biofilm processes: review of microfluidic methods. *Lab on a chip*. 2015; 15(1): 23–42. <https://doi.org/10.1039/c4lc01095g> PMID: 25385289
117. Giaouris E, Heir E, Hébraud M, Chorianopoulos N, Langsrud S, Mørretrø T, et al. Attachment and biofilm formation by foodborne bacteria in meat processing environments: causes, implications, role of bacterial interactions and control by alternative novel methods. *Meat Sci*. 2014; 97(3):298–309. <https://doi.org/10.1016/j.meatsci.2013.05.023> PMID: 23747091

118. Brasil. Ministério da Saúde. Manual Técnico de Diagnóstico Laboratorial da Salmonella spp. 2011. [Cited 2021 September 9] Available from: [https://bvsmms.saude.gov.br/bvs/publicacoes/manual\\_tecnico\\_diagnostico\\_laboratorial\\_salmonella\\_spp.pdf](https://bvsmms.saude.gov.br/bvs/publicacoes/manual_tecnico_diagnostico_laboratorial_salmonella_spp.pdf).
119. Garcia KCO, Corrêa IMO, Pereira LQ, Silva TM, Mioni MS, Izidoro ACM, et al. Bacteriophage use to control Salmonella biofilm on surfaces present in chicken slaughterhouses. *Poult Sci.* 2017; 96(9):3392–3398. <https://doi.org/10.3382/ps/pex124> PMID: 28595324
120. Sereno MJ, Ziech RE, Druziani JT, Pereira JG, Bersot LS. Antimicrobial susceptibility and biofilm production by Salmonella sp. strains isolated from frozen poultry carcasses. *Rev. Bras. Cienc. Avic.* 2017; 19(1):103–108.
121. World Health Organization (WHO). Background document: the diagnosis, treatment and prevention of typhoid fever. 2003. [Cited 2021 October 25] Available from: [https://www.who.int/selection\\_medicines/committees/expert/22/applications/s6.2\\_typhoid-fever.pdf](https://www.who.int/selection_medicines/committees/expert/22/applications/s6.2_typhoid-fever.pdf).
122. Brasil. Ministério da Saúde, Secretaria de Vigilância em Saúde, Departamento de Vigilância Epidemiológica. Manual integrado de vigilância e controle da febre tifóide. 2008. [Cited 2021 October 1] Available from: [https://bvsmms.saude.gov.br/bvs/publicacoes/manual\\_vigilancia\\_controle\\_febre\\_tifoide.pdf](https://bvsmms.saude.gov.br/bvs/publicacoes/manual_vigilancia_controle_febre_tifoide.pdf).